

A Case of Hepatocellular Carcinoma Metastasizing to the Palatine Tonsil

Kei Nakahara, Satoshi Matsuo, Hiromi Takeuchi and Hisaaki Ikoma

Department of Oto-Rhino-Laryngology, Faculty of Medicine, Tottori University, Yonago 683-0826, Japan

This report presents a very rare case of a patient with a metastasis in the left palatine tonsil from hepatocellular carcinoma (HCC). A 69-year-old man suffered from a mass of about 20 mm in the left palatine tonsil. The biopsy revealed that the mass in the palatine tonsil was a metastatic lesion from HCC. The patient underwent a left palatine tonsillectomy, and metastasis from HCC has not recurred since the operation. With the presentation of this case, we have looked through the literature reported so far.

Key words: tonsillar metastasis; hepatocellular carcinoma; first manifestation

Metastases from hepatocellular carcinoma (HCC) to the oral cavity are rare. We have recently experienced the case of a patient with tonsillar metastases from HCC, which was the first manifestation of the disease. The present article describes the case, with a thorough review of the literature.

Patient Report

A 69-year-old man with a mass of about 20 mm in the left palatine tonsil visited the Department of Oto-Rhino-Laryngology Clinic, Tottori University Hospital in March, 1999. The patient had noticed this mass for about 8 months. During this period, the patient removed it by himself with no complication, but it recurred immediately. The mass existed in the upper portion of the left palatine tonsil (Fig. 1), and its surface was necrotic. Magnetic resonance imaging showed that it was localized in the left palatine tonsil, without invasion to the ambient organs (Fig. 2). Pathological findings were metastases from HCC (Fig. 3). Although he had suffered from alcoholic liver disease with fatty liver for 10 years, there had been no evidence of HCC up to that moment; the tonsillar mass was the first demonstration of the tumor.

Clinical examinations revealed that a tumor

70 × 80 mm in size existed in the right hepatic lobe, which invaded the portal vein. There were no other metastases other than the left palatine tonsil, the alpha-fetoprotein level was positive (1216.0 ng/mL) and hepatic disorders were not severe (Child-A). The disease was non-curable.

The patient underwent a left palatine tonsillectomy under general anesthesia for removing the pharyngeal discomfort and for prevent-



Fig. 1. Oral cavity of the present patient. There is a tumor in the upper portion of the left palatine tonsil.

Abbreviation: HCC, hepatocellular carcinoma




Fig. 2. Magnetic resonance images (T2-weighted coronal views). The tumor is localized in the left palatine tonsil.

ing dyspnea, dysphagia and severe bleeding. The tonsil had no adhesion to the surrounding tissue, and there was no great loss of blood. No evident invasion could be found during the operation. We could not find any evidence of recurrence up to 4 months after the operation.

Discussion

Metastases to the tonsil are uncommon. The known primary tumors are malignant melanoma, carcinomas of the breast, lung and stomach, hypernephromas and seminomas (Brownson et al., 1979). Rarely observed cases are metastases from renal carcinoma, pancreatic carcinoma, Wilms' tumor, choriocarcinoma and mixed mesodermal tumor of the uterus (Fernández Aceñero et al., 1996). Tonsillar metastasis from HCC is quite rare. In Japan, at least 7 Japanese patients were reported from 1981 to 1994. In the West, we could find only 1 patient with HCC metastasizing to the palatine tonsil, reported from San Carlos University Hospital in Spain in 1996 (Llanes et al., 1996).

The primary tumors of tonsillar metastases reported in Japan (Table 1) are: carcinomas of

the lung, stomach, liver, colon, kidney, breast, pancreas, thyroid gland and gallbladder; malignant melanoma; choriocarcinoma; seminoma and mesothelioma (Funai et al., 1981; Mori et al., 1987; Kadowaki and Ikoma, 1990; Saiki et al., 1992; Kawai et al., 1993; Mochimatsu et al., 1993; Akiyama et al., 1995; Hando et al., 1995; Yamauchi et al., 1995). On the other hand, the reported metastatic sites in the head and neck from HCC are the orbit, paranasal sinus, gingivobuccal mucosa, palatine tonsil and lingual tonsil (Saiki et al., 1992; Kawai et al., 1993).

Why are tonsillar metastases rare? It has been hypothesized that translymphangial metastasis is rare to occur because the tonsil has no afferent lymph duct (Draizin et al., 1978). The tonsil has a high ability to eliminate cancer cells by immunosurveillance (Funai et al., 1981), so tonsillar metastases are rarely searched for at general postmortem examinations (Tanaka et al., 1981).

The most common sites for extrahepatic metastases from HCC are the lung, peritoneum, bone, intracelical organs and adrenal gland. The patient's tumor must have metastasized through the tonsillar hilus because HCC usually causes transvascular metastasis.