

Inflammatory Pseudotumor of the Lung Identified by ¹⁸F-Fluorodeoxyglucose Positron Emission Tomography: A Patient Report

Yuzo Takagi, Hiroshige Nakamura*, Ken Miwa*, Yoshin Adachi*, Shinji Fujioka*, Tomohiro Haruki*, Yuji Taniguchi* and Yasushi Horie†

*Center for Clinical Residency Program, *Division of General Thoracic Surgery and †Pathology Division, Tottori University Hospital, Yonago 683-8504 Japan*

The patient was a 71-year-old man with an abnormal shadow on chest X-ray. Computed tomography (CT) of the chest showed a tumor of 38 mm in size in the upper lobe of the left lung S³. A CT-guided lung biopsy was performed, but no malignancy was observed. ¹⁸F-fluorodeoxyglucose (FDG) positron emission tomography (PET) revealed a high FDG uptake in the tumor: the maximum and mean values of the standard FDG uptake in the early phase were 7.3 and 5.3, respectively, and those in the latter phase, 8.3 and 5.9, respectively. The hilar and mediastinal lymph nodes also showed positive for high FDG uptakes. We strongly suspected lung cancer, and performed a left upper lobectomy by video-assisted thoracic surgery. The pathological diagnosis was an inflammatory pseudotumor of the lung: it is a rare disease but often requires differentiation from lung cancer. Literature has been few on FDG-PET about inflammatory pseudotumor of the lung. Differentiation of the disease from lung cancer was especially difficult in the present patient, because both lymph nodes and the tumor showed high FDG uptakes.

Key words: ¹⁸F-fluorodeoxyglucose positron emission tomography; lung inflammatory pseudotumor; video-assisted thoracic surgery

Inflammatory pseudotumor of the lung has been more frequently detected due to the recent progress in diagnostic imaging, but it is difficult to differentiate from lung cancer. In many patients, the diagnosis is confirmed with surgery. We have encountered a male patient highly suspected of lung cancer due to his high uptake of ¹⁸F-fluorodeoxyglucose (FDG) in FDG-positron emission tomography (PET). We diagnosed his disease as inflammatory pseudotumor of the lung via video-assisted thoracic surgery. Few literature focuses on FDG-PET and inflammatory pseudotumor of the lung. The present patient showed a high FDG uptake also in the lymph node, which made us confused in differentiating from lung cancer. We herein present the case

of our patient, with bibliographical considerations as well.

Patient Report

This 71-year-old man had undergone medical examinations every year, but no abnormalities were detected. In July 2007, he had general fatigue and hemoptum, and visited nearby hospital where an abnormal shadow in the left upper lung was found on examination. He was referred to our University Hospital. Laboratory findings at the time of admission were a white blood cell count of 6,100/ μ L without any left shift of the nucleus, and the

Abbreviations: CT, computed tomography; FDG, ¹⁸F-fluorodeoxyglucose; PET, positron emission tomography

C-reactive protein level of 0.52 mg/dL. General or biochemical examinations of the blood showed no other abnormalities, and all of the tumor markers were negative. Chest X-ray showed a nodular density with approximately 40 mm in size in the left upper lung (Fig. 1). Chest computed tomography (CT) visualized a 38 × 32 mm tumor in size having broad contact with the parietal pleura in the left upper lobe S³ (Fig. 2). The periphery of the tumor was irregular, and the boundary was relatively well defined. Lymph nodes were enlarged at both the hilar and the mediastinal regions. In FDG-PET (Fig. 3), we observed a high FDG uptake just matching the tumor. The maximum and mean values of the standard FDG uptake in the early phase were 7.3 and 5.3, respectively, and those in the latter phase, 8.3 and 5.9, respectively. In addition, high FDG uptake was also observed in both the hilar and the mediastinal regions, so lung




Fig. 2. Chest CT revealed an irregular tumor, 38 × 32 mm in size, having broad contact with the parietal pleura in the left upper lobe S³. CT, computed tomography.




Fig. 3. FDG-PET revealed a high FDG uptake just matching the tumor, the hilar and mediastinal lymph nodes (arrows). FDG, ¹⁸F-fluorodeoxyglucose; PET, positron emission tomography.



Fig. 1. Chest X-ray showed an irregular-circumscribed abnormal shadow in the left middle field of the lung.

cancer and metastases to the lymph nodes were strongly suspected. The CT-guided lung biopsy resulted in nonspecific inflammatory findings, and no malignant cells were observed. However, on considering FDG-PET findings, lung cancer could not be ruled out. Therefore, thoracoscopic surgery was performed after obtaining informed consent from the patient. The tumor had firmly adhered to the parietal pleura, and a needle biopsy exhibited no malignant cells on pathological tissues frozen intraoperatively. We suspected an inflammatory pseudotumor, and performed a simple left upper lobectomy by video-assisted thoracic surgery. The resected tumor (Fig. 4) was 38 × 32 mm in size, sectionally xanthochromatic with mild anthracosis. Its shape was relatively regular in the periphery, well defined, solid and elastically soft. In the pathological tissue (Fig. 5), we found a lesion that was mainly comprised of organized pneumonic fibrosis, with inflammatory cell infiltration involving lymphocytes and plasma cells. When the image of the area was enlarged on the monitor, we observed inflammatory cells, fibrosis and spindle-shaped mesenchymal cells, which led us to the diagnosis of inflammatory pseudotumor, fibrous histiocytic type of the lung. The patient's postoperative course was good, and no recurrence has been observed.